


Chemistry B Final Exam Practice Quiz

Name: _____


1. Find the electronegativity difference between carbon (C) and chlorine (Cl), and indicate the type of bond that will form.
- 0.0 – 0.4 nonpolar covalent
 - 0.4 – 1.0 moderately polar covalent
 - 1.0 – 2.0 very polar covalent
 - ≥ 2 ionic

2. Which measurement correctly reflects the volume shown in this graduated cylinder?



- 36 mL
- 36.2 mL
- 36.20 mL
- 36.200 mL

3. Mercury forms a convex meniscus in glass, as shown in the picture to the right. This indicates that _____
- There are no intermolecular forces present.
 - The adhesive forces are stronger than the cohesive forces.
 - The cohesive forces are stronger than the adhesive forces.



4. What is the percent composition of CO₂?
- 27.29 % C and 72.71 % O
 - 27.29 % C and 36.36 % O
 - 42.88 % C and 57.12 % O
 - 42.88 % C and 72.71 % O

5. Determine the empirical formula of a compound with 87.4 % nitrogen and 12.6 % hydrogen.

- NH
- NH₂
- N₂H₄
- N₆H₁₂

6. Which of the following is an empirical formula?


- C₆H₁₂
- C₄H₈
- C₃H₈
- C₃H₆

7. How many milliliters of 6.00 M HCl are needed to make 0.250 L of 3.00 M solution?
- 0.125 mL
 - 0.50 mL
 - 125 mL
 - 500 mL

8. A 1.50 L balloon is sealed at 25 °C. Predict the volume if the balloon is heated to 35 °C.

- 1.07 L
- 1.45 L
- 1.55 L
- 2.10 L

9. How many moles of hydrogen gas are required to produce 4 moles of NH₃?



- 6 mol H₂
- 4 mol H₂
- 2 mol H₂
- 1 mol H₂

10. During an experiment you calculate that you should get 5.4 g of product from a reaction, but you only obtain 3.9 g of the product in the lab. What is your percent yield for this reaction?

- 1.4 %
- 7.2 %
- 28 %
- 72 %

11. Which of the following must be true of a reaction, if the change in heat involved in a chemical reaction has a positive sign?



- Heat is lost to the surroundings
- Heat is gained from the surroundings
- No heat is exchanged during the reaction

12. Which of the following needs the greatest amount of heat to raise the temperature 1°C?

- 1 g Aluminum, $C_p = 0.90 \frac{\text{J}}{\text{g}^\circ\text{C}}$
- 1 g Iron, $C_p = 0.46 \frac{\text{J}}{\text{g}^\circ\text{C}}$
- 1 g Silver, $C_p = 0.24 \frac{\text{J}}{\text{g}^\circ\text{C}}$
- 1 g Platinum, $C_p = 0.13 \frac{\text{J}}{\text{g}^\circ\text{C}}$

For each of the following problems involving calculations, show all work including units. If appropriate, indicate the equation used. Be sure to complete any conversions necessary. Record your answer with units in the box.

13. Determine the mass of nitrogen trihydride gas (NH₃) produced when 34.1 L of nitrogen gas, at STP, reacts with excess hydrogen.



14. Complete the table.

Draw the Electron Dot Structure	Draw 3-D structure & Name VSEPR Shape		Check(✓) all forces present & Circle or box the <input checked="" type="checkbox"/> to identify the strongest.	
		Indicate Polarity		
CH₃F	3-D Structure:	Shape Name: Polar or Nonpolar?	dispersion	<input type="checkbox"/>
			dipole-dipole	<input type="checkbox"/>
			hydrogen bonding	<input type="checkbox"/>
NH₃	3-D Structure:	Shape Name: Polar or Nonpolar?	dispersion	<input type="checkbox"/>
			dipole-dipole	<input type="checkbox"/>
			hydrogen bonding	<input type="checkbox"/>
SF₆	3-D Structure:	Shape Name: Polar or Nonpolar?	dispersion	<input type="checkbox"/>
			dipole-dipole	<input type="checkbox"/>
			hydrogen bonding	<input type="checkbox"/>
15. Based on the types of intermolecular forces you identified, which of the molecules in the table above is likely to adhere most strongly to water? (circle one) CH ₃ F NH ₃ SF ₆		16. Which of the molecules in the table above is likely to be the most volatile (evaporate the easiest)? (circle one) CH ₃ F NH ₃ SF ₆		

17. What is the volume of a sealed container with 12.7 grams of O₂ gas at 45°C and 2.5 atm?

$$C_{solid} = 2.10 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} \quad C_{liquid} = 4.18 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} \quad C_{gas} = 2.00 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} \quad \Delta H_{fus} = 334 \frac{\text{J}}{\text{g}} \text{ for H}_2\text{O} \quad \Delta H_{vap} = 2260 \frac{\text{J}}{\text{g}} \text{ for H}_2\text{O}$$

18. Calculate the change in energy (in kJ) needed to convert 25.0 g of steam at 100.0 °C to ice at -15.0 °C by following steps a and b below.

a. Using only variables, write the equation you will use to calculate the change in energy for the process described above.

b. Complete the calculation and report the total change in energy in kJ.

